



Mid-rapidity K_0 and Λ anisotropy parameters from Au+Au collisions at RHIC


Jinghua Fu
(for STAR collaboration)
IOPP/LBNL

1. Motivations
2. Theoretical predictions
3. Analysis method
4. Result

Azimuthal anisotropy v_2

Sensitive to initial/final conditions and equation of state (EOS) !

coordinate-space-anisotropy \leftrightarrow momentum-space-anisotropy




$$\varepsilon = \frac{\langle y^2 - x^2 \rangle}{\langle y^2 + x^2 \rangle}$$

$$v_2 = \langle \cos 2\varphi \rangle, \quad \varphi = \tan^{-1}\left(\frac{p_y}{p_x}\right)$$



Relativistic Hydrodynamics





Low Pt mass dependence

P.Huovinen, P.F.Kolb, U.Heinz, P.V.Ruuskanen,
S.A.Voloshin Physics Letters B503(2001)58-64




High Pt Azimuthal Asymmetry



M. Gyulassy, I. Vitev and X.N. Wang,
Phys. Rev. Lett. 86, 2537(2001).



Parton cascade & string melting




Zi-wei Lin and C.M. Ko,
Phys.Rev. C65(2002), pp034904



- Λ and K_s
- ★ Less resonance decay contributions
 - ★ Smaller hadronic cross section ----- Sensitive to partonic interaction
 - ★ PID up to high p_t ($\sim 3\text{GeV}$) ($\sim 6\text{GeV}$)

Reaction Plane Angle Distribution



Event and track selection


$|V_z| < 75$ cm
 $0.1 < P_t < 2$ GeV/c
 $|\eta| < 1.0$

global dca cut
 K_s v2: $gDca < 1$ cm
 Λ v2: $gDca < 1$ cm and no proton like track

201K minbias events





Reaction Plane Resolution



**Maximum resolution : 0.68 for $K_s v2$
0.65 for $\Lambda v2$**




V0 Vertex Reconstruction






K_0 and Λ identifications




STAR Preliminary



Background subtraction



**Subtract background
in each pt and ϕ bin**





**ϕ distribution after
background subtraction**



$$V_2 = \langle \cos[2(\phi - \Psi_R)] \rangle$$




Phase Space Coverage



-1 < y < 1 and 0.2 < Pt < 2 GeV/c




Centrality dependence of $v_2(p_T)$ at 130GeV



STAR Preliminary




$v_2(p_T, m)$ for Λ and K_S^0 at 130GeV





Azimuthal anisotropy of K_0 , Λ



STAR Preliminary




Summary

- 1) Event anisotropy v_2 of K_0 and Λ increase as a function of transverse momentum within $P_T \leq 1.5\text{GeV}/c$ and seem to be saturating at higher momentum
- 2) Stronger P_T dependence is observed in more peripheral collisions
- 3) The integrated values of V_2 show a mass dependence, hydrodynamic model results are not inconsistent with our measurement





$V_2(p_T, m)$ from Minbias Events



STAR Preliminary



$V_2(P_t)$ from 200Gev & 130Gev



STAR Preliminary